

# LIS / OTD Gridded Products

**V 2.2 : 1995 - 2005**

**Documentation and Examples**

**1 September 2006**

# Gridded vs Orbit products

— [ For most climatological or time series studies, the gridded products are preferable to the individual orbit HDF's

— [ They apply the MSFC best-estimate of spatially and diurnally varying detection efficiency, and correct for instrument viewing time.

— [ OTD and LIS datasets have been intercalibrated and merged for the products. Nonetheless, care should be used in interpreting time series results between the OTD-only window (5/95-11/97), the overlap window (12/97-4/00) and the LIS-only window (12/97-present).

— [ Orbit HDF's should be used if individual flash times, locations, properties, etc are needed (e.g., for cross-sensor comparisons)

# Naming Convention

— [ **HR = High Resolution (0.5 degree grid)**

— [ **LR = Low Resolution (2.5 degree grid)**

— [ **FC = Annualized Climatology (full year)**

— [ **SC = Seasonal Climatology (3-monthly)**

— [ **MC = Monthly Climatology**

— [ **AC = Annual Climatology (daily)**

— [ **DC = Diurnal Climatology (hourly)**

— [ **TS = Time Series (daily or monthly)**

# Master HDF Dataset List

	Annualized	Time Series	Seasonal	Monthly	Daily	Hourly
HRFC	•					
LRFC	•					
HRMC	•		•	•		
HRAC	•				•	
LRAC	•				•	
LRADC	•				•	•
LRDC	•					•
LRMTS		•		•		
LRTS		•			•	
LRACTS	•				•	



# Smoothing

— [ **Note:** The nominal grid resolution may be higher than the resolution of the data going into it. Significant spatial and temporal smoothing may have been used to generate meaningful estimates. This is documented for each individual file.

— [ E.g., the HRAC (High Resolution Annual Climatology) contains data at daily resolution (365 grids), but each grid represents a +/- 55-day moving average.

# Dates of Data

- [**OTD:** May 1995 - April 2000

- Global coverage

- $\sim 60\%$  uptime; 110-day diurnal alias window

- [**LIS:** Jan 1998 - Dec 2005

- Tropical ( $\sim 35\text{N/S}$ ) coverage. Slightly more after TRMM orbit boost.

- $\sim 99\%$  uptime; 98-day diurnal alias window

- Climatology grids are full years; time series may not be aligned with year boundaries

# File Type / Contents

- Files are HDF Version 4

- Simple SDS (Scientific Data Set) Grids, 2-or-more dimensional

- Accessible with NCSA HDF tools, many 3rd-party applications, or the included IDL "getgrid.pro" procedure

- Units are embedded with grid dimensions

# Key grid within each file

— [ **XXXX\_COM\_FR**

- **COM**bined **F**lash **R**ate density (flashes / area / time)
- Merged estimate from OTD and/or LIS
- Scaled by best estimate of diurnally-varying detection efficiency (DE)
- DE-Scaled Flashes divided by Viewtime

# Diagnostic grids within some files

— [ **XXXX\_OTD\_FR, XXXX\_LIS\_FR**

— Uncombined **OTD** and **LIS** Flash Rate Densities

— [ **XXXX\_XXX\_VT**

— Total **ViewTime** used to compute FR

— [ **XXXX\_XXX\_RF, XXXX\_XXX\_SF**

— Total **Raw, DE-Scaled Flash** counts used to compute FR




# HRFC (High Resolution Full Climatology)

— [ SDS: HRFC\_COM\_FR:

— 0.5 deg resolution, 720x360; flashes / km<sup>2</sup> / year


\*Also:

HRFC\_OTD\_FR,  
HRFC\_LIS\_FR,  
HRFC\_OTD\_SF,  
HRFC\_OTD\_RF,  
HRFC\_OTD\_VT,  
HRFC\_LIS\_SF,  
HRFC\_LIS\_RF,  
HRFC\_LIS\_VT



# HRFC (High Resolution Full Climatology)

Total viewing time for HRFC\_COM\_VT (hours):



# LRFC (Low Resolution Full Climatology)

— [ SDS: LRFC\_COM\_FR:

— 2.5 deg resolution, 144x72; flashes / km<sup>2</sup> / year

\*Also:

LRFC\_OTD\_FR, LRFC\_LIS\_FR, LRFC\_OTD\_SF, LRFC\_OTD\_RF, LRFC\_OTD\_VT,  
LRFC\_LIS\_SF, LRFC\_LIS\_RF, LRFC\_LIS\_VT




# HRMC (High Resolution Monthly Climatology)

- [ SDS: HRMC\_COM\_FR (monthly):
  - 0.5 deg resolution, 12x720x360; flashes / km<sup>2</sup> / day
  - 30-day average centered on 15th of each month
  - 2.5 deg moving average applied to each 0.5 deg grid
- [ SDS: HRSC\_COM\_FR (seasonal):
  - As above, but:
  - 90-day average centered on 15th of each month



# HRMC (High Res Monthly Climatology Grids)


HRMC





# HRMC (High Res Seasonal Climatology Grids)

HRSC






# HRAC (High Resolution Annual Climatology)

- [ SDS: HRAC\_COM\_FR (daily):
  - 0.5 deg resolution, 365x720x360; flashes / km<sup>2</sup> / day
  - 110-day moving average centered on each day of year
  - 2.5-deg moving average applied to each 0.5 deg grid
  - Minimal diurnal aliasing, smooth annual cycle evolution


# HRAC (High Resolution Annual Climatology)

Sample animation:




# HRAC (High Resolution Annual Climatology)

Sample animation






# HRAC Peak Lightning Months






# HRAC Seasonal Cycle Amplitude





# HRAC Peak Seasonal Flash Rate



# LRAC (Low Resolution Annual Climatology)

- [ SDS: LRAC\_COM\_FR (daily):
  - 2.5 deg resolution, 365x144x72; flashes / km<sup>2</sup> / day
  - Low resolution version of HRAC



# LRADC (Low Resolution Annual / Diurnal Climatology)

## — [ SDS: LRADC\_COM\_SMFR:

- 2.5 deg resolution, 365x6x144x72; flashes / km<sup>2</sup> / day
- Smoothing as in HRAC, LRAC
- 6x 4-hourly UTC bins (0-4, 4-8, ... 20-24)


## — [ SDS: LRAD\_COM\_SMFR2:

- 2.5 deg resolution, 365x12x144x72; flashes/ km<sup>2</sup> / day
- 12x 2-hourly UTC bins (0-2, 2-4, ... 22-24)


# LRDC (Low Resolution Diurnal Climatology)

— [ SDS: LRDC\_COM\_FR (daily):

- 2.5 deg resolution, 24x144x72; flashes / km<sup>2</sup> / hour
- 24 local hour bins (0-1, 1-2, ... 23-24 LST)



Local



Converted to UTC

# LRMTS (Low Resolution Monthly Time Series)

— [ SDS: LRMTS\_COM\_FR:


— 2.5 deg resolution, 144x144x72; flashes / km<sup>2</sup> / day

— 144 months beginning 1/1/05; first 5 and last 8 months empty

— 7.5 deg spatial moving average

— 110-day or 98-day temporal smoothing (OTD, LIS)

— OTD and LIS grids weighted by respective viewtimes during overlap window (12/97 - 04/00)




# LRTS (Low Resolution Time Series)

— [ SDS: LRTS\_COM\_FR:

- 2.5 deg resolution, 3653x144x72; flashes / km<sup>2</sup> / day
- 4261 days beginning 1/1/05; first and last few months empty
- 7.5 deg spatial moving average
- 110-day temporal smoothing

— [ SDS: LRTS\_LIS\_FR:

- As above, but LIS-only; 98-day smoothing



# LRACTS (Low Res Annual Clim Time Series)

- This is a special version of the LRAC (2.5 degree annual climatology) created for use with the LRTS time series product
- It uses the same spatial / temporal smoothing as LRTS
- It can thus be used as a “baseline” against which time series anomalies in the LRTS can be compared.
- This is more accurate than comparing against the LRAC




# LRTS / LRACTS

## Example

Full time series  
for a (7.5x7.5  
smoothed) location  
centered at 61W,  
2S (near TRMM-  
LBA / Rondonia)

LRTS is the solid  
curve; LRACTS  
(repeated  
annually) is the  
dashed curve





# getgrid.pro

— [ IDL Routine (v5 or higher)

— [ *getgrid*, **hdfname**, **sdsname**, gridvar, dims, dimnames, dim0, dim1, dim2

— hdfname: HDF file name (full path if not local); e.g. "LISOTD\_HRAC.hdf"

— sdsname: SDS grid name, e.g. "HRAC\_COM\_FR"

— gridvar: IDL variable in which the grid will be returned

— dims, dimnames, ...: IDL variables in which the dimensions will be returned